

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.

Do not take more than the prescribed dose.

Overdose can lead to serious adverse effects and even death. Symptoms of overdose include sore throat, fever, mouth ulcers, diarrhoea, vomiting, skin rashes, bleeding or unusual weakness. If you think you have taken more **methotrexate** than prescribed, contact a doctor straight away.

Always show this card to healthcare professionals not familiar with your **methotrexate** treatment to alert them about your once weekly dose eg if you go to hospital or have a change in care.

For more information, please read the patient leaflet in your medicine pack.